

ICHTHYOFAUNA AND FISH PRODUCTION IN THE POTIASOLA WETLAND OF THE BRAHMAPUTRA BASIN

R. BORDOLOI

Department of Zoology, D. R. College, Golaghat, Assam, India

ABSTRACT

Investigation on the wetlands was conducted from January 2008 to December 2009. A total of 39 species have been recorded and belonging to 17 families during the reporting period. Out of these species 37 species are indigenous species having ornamental values, edible as well as medicinal values while two species are exotic i.e., *Cyprinus carpio* (Lin.) & *Ctenopharyngodon idella*. The wetland was dominated by live fishes and among them, *Clarias batrachus* is a dominant species with 44% and followed by *Channa sp.* with 35% and *Anabas testudineus* with 21%. As per IUCN, out of the recorded species in the beel, 2.56% species as Endangered; 10.25% species as Vulnerable (VU); 7.69% species as Lower Risk least concern (LRlc), 38.46% species Near threatened (LRnt), 35.89 species are still Not assessed (NA) and 3% species are exotic. The rate of fish production from the beel has been estimated to be an average of 600 kg /yr. (25.65 kg/h).

KEYWORDS: Closed Wetland, Potiasola, Beel

INTRODUCTION

Wetlands (beels) are highly productive of the ecosystem. The North Eastern region of India is considered as one of the hot spots of freshwater fish biodiversity in the world (Kottelat & Whitten, 1996). The mixing of drainages and their fish fauna in the geographical past has rendered the Assam Himalayas as very important from faunistic point of view. Most significant contributions on fish diversity and beel ecology are those of Agarwala (1996) in Tamrange wetland; Biswas & Boruah (2000) in lentic and lotic water bodies of upper Brahmaputra basin; Dutta (2002) in “closed” and “open” beels in upper Assam; Bera et al. (2008) in Deepor beel; Singh *et al.*, (2009); Dakua et al., (2009); Abujam *et al.* (2011 & 2012) in Maijan beel; Hussain & Biswas (2011) in wetland of Dhemaaji; Paswan *et al.* (2012) in Borsala beel of Jorhat. So far 267 fish species belonging to 114 genera under 38 families and 10 orders has been recorded and reported from the region.

This is about 33.13% of total Indian fresh water fishes (Sen, 2000). Sarkar and Poonia (2000) evaluated the ornamental value of 172 fish species occurring in North Eastern Region. As per NBFGR (2002) report among the states, Assam has the largest number of ichthy species (200), followed by Arunachal Pradesh (169), Meghalaya (165), Tripura (134), Manipur (121), Nagaland (68) and Mizoram (48). The status and export potentialities of indigenous ornamental fishes of India were highlighted by some workers (Dey, 1980 & 1982), Sen and Dey (1984), Nath (1986 & 1987) and Tamang (1992). The ornamental fish diversity of the North eastern region of India offers potentialities for its trade in global market. But aquatic fish species are highly threatened species in the North Eastern region as well as in the world due to anthropogenic pressure. Considering the above view, the present study has taken up the fish diversity and its production from Potiasola beel.

MATERIAL AND METHODS

The Potiasola wetland (closed beel) located in the geographical ordinates of 26°48′-26°49′N and 94°08′-94°10′ E. and situated at about 5 km from North east of Jorhat town, Assam. The present investigation was carried out in the beel from January 2008 to December 2009. Fish samples were collected for quarterly basis at the time of catch at the landing site of the *beel*. The fish species were identified with the help of standard procedure of Talwar and Jhingran (1991) and Jayaram (1999). Fish samples were preserved in 5% formalin for further investigation. The individual species was weighed and recorded after collection. Fish yield are estimated by direct observations on catch of individual fishermen (Jhingran and Dutta, 1968) and also with the help of local fishermen and Mohalder having more than 20 years fishing experiences.

RESULTS

Hydro topographic measurement of the Potiasola wetlands is given in Table 1. The area increases up to about 20.00 ha. during rainy season, due to the surrounding low-lying paddy fields, which become an integral part of the beel. Depth increases up to about 6 meters at the season. During dry season, the area decreases up to about 10.00 ha. Depth decreases up to half meter. Altogether 39 species belonging 17 families have been encountered during study period (Table 2). Among the family Cyprinidae was found to be highest species (12) and followed by Belontiidae and Channidae with 4 species each and Bagridae, Chandidae, Mastacembelidae, Nandiidae and Siluridae with 2 species each. Some of the indigenous species having ornamental values, edible as well as medicinal values while two species are exotic i.e., *Cyprinus carpio carpio* (Lin.) & *Ctenopharyngodon idella*.

As per IUCN (Table 2), out of the recorded species in the *beel*, 2.56% species as Endangered; 10.25% species as Vulnerable (VU) (facing a very high risk of extinction in the wild); 7.69% species as Lower Risk least concern (LRlc), 38.46% species Near threatened (LRnt), 35.89 species are still Not assessed (NA) and 3% species are exotic. The wetland was dominated by live fishes and among them, *Clarias batrachus* is a dominant species with 44% and followed by *Channa sp.* with 35% and *Anabas testudineus* with 21% (Table 3 & 4). Total production was 710 kg during the study period. Out of that 37% production (Table 3) from catfishes followed by murrels (29%), miscellaneous (18%) including *Anabus testudineus*, feather back (8%), carps (5 %) and exotic sp. (3 %).

DISCUSSIONS

Wetlands are the creation of the river Brahmaputra in the flood plain area. Water level of the Potiasola beel is dependent upon the rainwater from surrounding paddy fields and connecting channel. Interestingly two exotic species were also encountered in the open beel. This investigation reveals that there has been drastic reduction in the abundance and distribution range of fishes in North East region due to habitat modification, overexploitation and other anthropogenic pressure (Deka *et al.* 2005). The International Union for Conservation of Nature and Natural Resources (IUCN) has categorized the conservation status of the recorded fish species. The highest percentage was found to be LRnt & NA category and followed by VU, LRlc, and EN category respectively (Figure 1). Out of that 37% production from catfishes followed murreles, miscellaneous including *Anabus testudineus*, feather back, carps and exotic sp. (Figure 2). The *beel* was dominated by live fishes than carps species and among them, *Clarias batrachus* is a dominant species and next followed by *Channa sp.* and *Anabas testudineus* (Figure 3). Similar record was observed by Bordoloi (2010). Family Cyprinidae have highest species and followed by Belontiidae and Channidae. Similar observation was reported by Singh (2009); Dakua (2009) and Abujam (2012). Most of them have ornamental, edible values as well as medicinal values. These fishes also

have high price and demand in the local market. So, these fishes should be conserved and develop breeding technique in-situ and ex-situ condition.

Fish Production

On the basis of questionnaire, annual fish yield of the wetland was estimated about 25.5 kg/ha. The Central Inland Fisheries Research Institute, Barrackpore has estimated yields varying from 14 to 488 kg/ha/year. Average yields of 17 *beels* in the Brahmaputra valley is 134 kg/ha/year, compared to 285 kg/ha in 6 *beels* of the Barak valley (Sugunan and Sinha, 2000). The average yield of 23 *beels* in Assam across the districts is 173 kg/ha (Sugunan and Sinha, 2000). The present finding was found to be low in comparison to the other findings. Ecological degradation is the most important factor for fish declining. The ecological degradation of the beels started with the arrival of the water hyacinth a century ago. Rampant growth of this fast – growing weed obstructs the penetration of sunlight, inhibiting plank tonic growth and contributing to eutrophication by slowing down water currents and depositing debris at the bottom. The second phase of enhanced eutrophication resulted from the construction of embankments along almost the entire length of the river Brahmaputra and many of its tributaries after the devastating earthquake of 1950. The final onslaught on the wetlands has been from human activities such as buffalo and cattle rearing, agriculture and horticulture, and over fishing. These have resulted in further siltation and damage to the micro flora and water quality. The injudicious use of pesticides in farming activities has resulted in the accumulation of residue through surface runoff, leading to the problem of biomagnification. A number of fish species, such as *Puntius jerdoni*, *Bagarius bagrius* and *Semiplotus semiplotus* are on the verge of extinction (Dubey and Ahmed 1950). The use of unauthorized mesh size and use of a wide range of non – selective fishing gears such as mosquito nets in this state indicates that most fishers do not comply with the existing fishery act and fishing regulations and are not concerned with possible overexploitation of the stocks of commercially important species.

Table 1: Hydrotopographic Measurement of the Closed Wetlands

Parameters	Measurement
Area at Full Storage Level (ha)	20
Area at Dead Storage Level (ha)	10.25
Total volume at FSL m3)	31729.62
Total volume at DSL m3)	15864.81
Maximum length at FSL (m)	1000
Maximum width at FSL (m)	450
Maximum length at DSL (m)	450
Maximum width at DSL (m)	210
Maximum depth at FSL (m)	6
Minimum depth at FSL (m)	0.25
Mean depth (Dm) at FSL (m)	4.54
Maximum depth at DSL (m)	2.10
Minimum depth at DSL (m)	0.20
Mean depth (Dm) at DSL (m)	0.25
Volume development (VD) at FSL)	3.96
Volume development (VD) at DSL	0.25

Table 2: Status of the Recorded Fish Species of the Potiasola Beel

Family	Species	Status(IUCN)	Economic Importance
1. Anabantidae	1. <i>Anabas testudineus</i> (Ham.)	VU	Edible, Medicinal value, Aquarium species
2. Belontiidae	1. <i>Trichogaster fasciata</i>	NA	Edible, Aquarium species
	2. <i>T. labiosa</i>	NA	Edible, Aquarium species

Table 2: Contd.,

	3. <i>T. lalia</i>	NA	Edible, Aquarium species
	4. <i>T. sota</i>	NA	Edible, Aquarium species
3. Belonidae	1. <i>Xenentodon cancila</i> (Ham.)	LR-nt	Edible, Aquarium species
4. Bagridae	1. <i>Mystus cavasius</i> (Ham.)	LR-nt	Edible, Aquarium species
	2. <i>M. tengra</i> (Ham.)	NA	Edible, Aquarium species
5. Chandidae	1. <i>Chanda nama</i> (Ham.)	NA	Edible, Aquarium species
	2. <i>Parambassis ranga</i> (Ham.)	NA	Edible, Aquarium species
6. Clariidae	1. <i>Clarius batrachus</i> (Linn.)	VU	Edible, Medicinal value Aquarium species
7. Cobitidae	1. <i>Botia dario</i> (Ham.)	NA	Edible, Aquarium species
8. Cyprinidae	1. <i>Amblypharyngodon mola</i> (Ham.)	R-lc	Edible, Aquarium species
	2. <i>Cyprinus carpio carpio</i> (Lin.)		Edible,
	3. <i>Chela apter</i> (Ham.)	NA	Edible, Aquarium species
	4. <i>Cirrhinus mrigala</i> (Ham.)	LR-nt	Edible
	5. <i>Labeo bata</i>	LR-nt	Edible
	6. <i>Labeo gonius</i> (Ham.)	LR-nt	Edible
	7. <i>Puntius chola</i> (Ham.)	VU	Edible, Aquarium species
	8. <i>Puntius gelius</i> (Ham.)	NAS	Aquarium species
	9. <i>P. sophore</i> (Ham.)	LR-nt	Edible, Aquarium species
	10. <i>P. ticto</i> (Ham.)	LR-nt	Edible Aquarium species
	11. <i>Rasbora daniconius</i> (ham.).	NA	Edible, Aquarium species
	12. <i>Ctenopharyngodon idella</i>	Exotic	Edible
9. Channidae	1. <i>Channa gachua</i> (Ham.-Bloch.)	NA	Edible, Aquarium species
	2. <i>C. marulius</i> (Ham.)	LR-nt	Edible, Aquarium species
	3. <i>C. punctatus</i> (Bloch.)	LR-nt	Edible, Aquarium species
	4. <i>C. striata</i> (Bloch.)	LR-lc	Edible, Aquarium species
10. Clupeidae	1. <i>Gudusia chapra</i> (Ham)	LR-lc	Edible Aquarium species
11. Heteropneustidae	1. <i>Heteropneustes fossilis</i> (Bloch)	VU	Edible, Medicinal value, Aquarium species
12. Mastacembelidae	1. <i>Mastacembelus armatus</i> (Lacepede)	LR-nt	Edible Aquarium species
	2. <i>Macrognathus pancalus</i> (Ham)	LR-nt	Edible Aquarium species
13. Nandidae	1. <i>Nandus nandus</i> (Ham)	LT-nt	Edible, Aquarium species
	2. <i>Badis badis</i>	NA	Edible, Aquarium species
14. Notopteridae	1. <i>Notopterus notopterus</i>	LR-nt	Edible Aquarium fish
15. Siluridae	1. <i>Ompok pabda</i> (Ham)	EN	Edible, Aquarium fish
	2. <i>Ompok pabo</i> (Ham)	NA	Edible, Aquarium fish
16. Synbranchidae	1. <i>Monopterus albus</i> (Ham)	LR-nt	Edible, Aquarium species
17. Tetraodontidae	1. <i>Tetraodon lineatus</i> (Ham)	LR-nt	Edible, Aquarium species

Table 3: Monthly Fish Catch Statistics (Kg) in Potiasola Wetlands

2008 -09	J	F	M	A	M	J	J	A	S	O	N	D	SUM	MEAN
CAT FISH	2	2	5	2	3	4	20	20	40	40	60	60	258	39.69
MURREL	5	9	6	8	20	10	10	20	30	20	40	30	208	32
FEATHERBACK	0	0	1	0	5	5	5	9	5	10	10	5	55	8.462
MISCELLANEOUS INCLUDING ANABUS SPECIES	1	5	6	8	10	10	10	10	10	20	20	20	130	20
CARP	0	0	0	0	3	5	2	10	5	2	5	5	37	5.692
Carp (exotic)	0	0	0	0	2	3	1	2	3	3	3	5	22	3.667
TOTAL													710	710

Table 4: Production of Live Fishes in the Potiasola Wetlands

Live Fishes	Production (Kg)	%
<i>Clarias batrachus</i> & <i>Heteropneustes fossilis</i>	258	44.02
<i>Anabus testudineus</i>	120	20.47
<i>Channa sp.</i>	208	35.49
Total	586	




Figure 1: Status of Fishers in the Potiasola Wetland




Figure 2: Fish Composition in the Potiasola Wetland




Figure 3: Percentage of Live Fishes in the Potiasola Wetlands during Study

ACKNOWLEDGEMENTS

Acknowledged the University Grant Commission (UGC) for providing financial assistance to carry out the research work. I offer my heartiest thank to Principal & Head, Department of Zoology, D. R. College, Golaghat, Assam, for providing me the facilities to complete the work. Special thanks to Department of fisheries, Assam, India for providing me important information of wetlands. I wish to express my gratitude to Prof. U. C. Goswami, Dept of Zoology, Gauhati University for his guidance during preparation the manuscript.

REFERENCES

1. Abujam, S. K. S., S. Dakua, B. Bakalial, A. K. Saikia, S. P. Biswas, P. Choudhury, (2011), Diversity of Plankton in Maijan Beel, Upper Assam. *Asian J. Exp. Biol. Sci.*, 2(4): 562-568.
2. Abujam, S. K. S., G. Paswan, S. Dakua, A. K. Saikia (2012): Faunal diversity and ecological status of Maijan beel (Wetland). *Fishing Chimes*, 32(8): 43-48.

3. Agarwala, N. K., (1996), Limnology and fish productivity of Tamranga wetland in Bongaigaon district of Assam (India) with special reference to some productivity indicator, PhD Thesis, Gauhati University, Assam, pp. x+200.
4. Biswas, S. P. and S. Boruah, (2000), Fisheries ecology of the North-Eastern Himalaya with special reference to the Brahmaputra River. *Ecol. Eng.*, 16: 39-50.
5. Dakua, S., A. S. K. Singh, P. Choudhury, and S. P. Biswas, (2009), A preliminary investigation on the fish and fisheries of Majan beel in upper Assam. *Indian J. Environ. & Ecoplan.*, 16 (1): 263-270.
6. Hussian, Md. and S. P. Biswas, (2011), Physico-chemical characteristics of a floodplain lake of Dhemaji in upper Assam. *Bull. Life. Sci.*, (16): 81-91.
7. Kottelat, M. and Whitten, T. (1996): Fresh water biodiversity in Asia with special reference to fish. *World Bank Tech. Paper*, 343: 17-22.
8. Singh, A. S. K., S. Dakua and S. P. Biswas, (2009), Physico-chemical parameters and fish enumeration of Majan beel (wetland) of upper Assam. *Geobios*, 36: 184-188.
9. Paswan, G., Singh, A. S. K., Biswas, S. P. and Dey, M., (2012), A Study of Certain Physico-Chemical Parameters of Borsola Beel (Wetland) of Jorhat. (Ed.) *Proc. National seminar on 'Current Issues of Conservation and Wise Use of Wetlands in North Eastern Region of India'*, 291-297 p., (2012)
10. Dutta, P. K., (2002), A comparative study of the hydrobiology and fisheries of open and closed beels in Jorhat district of Assam. Ph. D. Thesis (unpublished), Dibrugarh University, India.
11. Bera, S. K., S. Dixit, S. K. Basumatary, and R. Gogoi, (2008), Evidence of biological degradation in sediments of Deepor beel Ramsar site, Assam as inferred by degraded Palynomorphs and fungal remains. *Current Science*, 95(2): 178-180.
12. Anon, 1997. Wetland of Assam. Assam Remote Sensing Application Centre, Guwahati. (Margate, G. R.; Obbert, M. Jr. and Wolf, C. L. : Glossary of Geology. American Geological Institute, Washington DC
13. Agarwala, H. 1994. Endangered sport fishes of Assam. *In: Threatened fishes of India*, Dehadrai, P.V., Das, P. and Verma, S.R. (eds.) *Nature Conservators*, Muzzafarnagar, 4: 209-212 pp.
14. Biswas, S. P. and Baruah, S. 2000. Problem and prospects for the development of fisheries in the Brahmaputra basin. *In: Fish Germplasm Inventory and Conservation in North East India*. Pooniah, A. G. and Sarkar. U.K. (eds.), NBFGR, Lucknow.
15. Bordoloi, R. 2006. Fish germplasm inventory and conservation on the wetland of Assam (abst.). pp – 43, National seminar on Zoology for human welfare, 2nd february, G.U. 2006.
16. Bordoloi, R. 2007. Conservation of Wetland habitats to increase abundance of fish diversity in the wetland of Assam. 22 April 2007, National seminar, a.s.c. pp- 19 (abst.)
17. Bordoloi, R. 2008. Problem and prospects of the wetland in the Jorhat district, Assam (abst.), pp-63, 14th & 15th ugc sponsored National seminar at, Gorgaon college, Assam.

18. Bordoloi, R. 2008. Fish and fisheries of a closed and open type wetland of the Jorhat district, Assam (abst.), PP- 207, National seminar on Biodiversity & Human Welfare, 29 – 31 Dec. 2008, G.U.
19. Bordoloi, R. 2008. Exploration of fish fauna, fish production and habitat conservation of Nahotia and Potiasola wetland of Jorhat district, Assam. pp. 150, proceedings of 53rd annual technical session, journal of a.s.c..
20. Bordoloi, R. 2010. Comparative study on fish and fisheries between a closed and an open type wetland of the Jorhat, District, Assam, Ph.D. Thesis, Gauhati University
21. Choudhury, M. 1987. An analysis of fish catch statistic in Dhir beel, Assam: Workshop on Development of beel of fishery in Assam, Assam Agri. Univ.: 47 -69 pp.
22. Dey, S. C. and Kar. D. 1987. Physico-chemical complexes of water and soil in Sone, an ichthyologic ally potential tectonic lake of Assam, J. Assam. Sc. Soc. 30(1) : 1-11
23. Dey, S.C. 1973: Studies on the distribution and taxonomy of the ichthyofauna of hill streams of Kamrup and Khasi-Garo regions of Assam with special reference to the functional morphology of some rheophilic fishes. Unpublished D.Sc Thesis, Calcutta Univ.
24. Dey, S.C. 1975: Ecomorphology of the ichthyofauna of the river Pagladia within Kamrup district, Assam. *J. Assam Sci. Soc.* Vol. 18(1): 1-5 pp.
25. Dey, S.C. 1976: a. Studies on the probable origin and distribution of the ichthyofauna of the river Brahmaputra, India, Rev. des. Trav, de 1. Inst. et. Tech. des. Poches Merit, Paris.
26. Dey, S. C. 1976: b. Significance of fishes in the hill streams of Assam and Meghalaya, India, Rev. des. de 1. Inst. et. Tech. des. Poches Merit, Paris.
27. Dey, S. C. 1976: c. *Semiplotus gangulyi*, a new *Cyprinid* from Pagladia river of Kamrup district of Assam, Rev. des. Trav, de 1. Inst. et. Tech. des. Poches Merit, Paris.
28. Dey, S. C. 1976 a: Studies on the probable origin and distribution of the ichthyofauna of the river Brahmaputra, India, Rev. des. Trav, de 1. Inst. et. Tech. des. Poches Merit, Paris.
29. Dey, S. C. 1976b: b. Significance of fishes in the hill streams of Assam and Meghalaya, India, Rev. des. de 1. Inst. et. Tech. des. Poches Merit, Paris.
30. Dey S. C. 1976 c: *Semiplotus gangulyi*, a new *Cyprinid* from Pagladia river of Kamrup district of Assam, Rev. des. Trav, de 1. Inst. et. Tech. des. Poches Merit, Paris.
31. Dey, S. C. and Lahon, B. 1980: Dial dynamics of rotifer biomass and species composition during winter in an oxbow lake of Assam, *Proc. Indian Sci. Cong.* 67 (3): 159pp.
32. Dey, S.C. 1982: A Critical Analysis on the fish and fisheries of Assam, *Proc. Nat. Seminar on Ichthyology.* 3:16.
33. Dey S. C. and Sen, N. 1982 Fish and Fisheries of some major drainages of Meghalaya, *Proc. All-Indian Sem. Ichthyology*, 3: 10-11 pp.
34. Dubey C.E. and Ahmed A. (1995). Problems for the conservation of freshwater fish metric resources in India and some possible solution. *NAGA.ICLARMQ.* 18(3):21-25.

35. Deka T.K 2005. Causes of fish depletion- A factor analysis approach, *NAGA, World Fish Centre Newsletter*, Vol. 28: No. 1 & 2, 37-42 pp.
36. Dutta Munshi, J. S. and Srivastava, M. P. 1988: Natural History of Fishes and Systematic of Freshwater Fishes of India, Narendra Publishing House, New Delhi.
37. Goswami, M. M. 1996. Ichthy-Ecological status of the beels of Assam with reference to their fish resources, *Projna Jour. GUTA*, vol.V:48-53.
38. Goswami, M. M., Deka, T. K., Singha, P. K., Sarma, P. K. and Kakati, M. 1999: Studies on some wetlands of Assam with reference to the Eutrophication stresses, *J. Ind. and Fish Soc. India*, 3(12).
39. Jhingran, V. G. and Dutta, P. 1968. The Inland fisheries resources of India. *Indian Soc. Agric. Stast.* 20© 68-78.
40. Jhingran, V. G. 1991. *Fish and Fisheries of India*, pp. 61 -67.
41. Jha, B. C. et al, (1968) :The Inland fisheries resources of India. *Indian Soc. Agric. Stast.* 20© 68-78.
42. Jayaram, K. C. 1999. *The Freshwater Fishes of the Indian Region*, Narendra Publ. House, Delhi, p. 551.
43. Kar, D. 1984. Limnology and fisheries of lake Sone in the Cachar district of Assam, India, PhD Thesis, Gauhati University, Assam pp. 201.
44. Kar, D. 2000a. Present status of fish biodiversity in South Assam and Tripura, In: Ponniah, A.G. and Sarkar, U.K. (Eds.) *Fish Biodiversity of N.E India*, NBFGR-NATP Pub. NO.2, p. 80-82.
45. Lahon, B. 1983. Limnology and fisheries of some commercial beels of Assam, India, PhD Thesis, Gauhati University (Assam), pp.351..
46. Nath, P. and Dey, S.C. 1990. *Fish and Fisheries of North East India*, vol. I (Arunachal Pradesh), pub I. A K Choudhury, Barrakpore, pp. 201.
47. Nath, S. 1986 The fish and fisheries of the Barak river system in Assam, India with special reference to the ecomorphology and distribution of its ichthyofauna, PhD Thesis, Gauhati University (Assam), pp. xii+350.
48. Nath, P. 1987: Systematics, Distribution and Ecology of Ichthyofauna of Arunachal Pradesh with particular reference to the Limnobiology of Itanagar Lake, PhD Thesis, Gauhati University, pp. 362.
49. Sen, T. K. 1985. The Fish fauna of Assam and the Neighboring North-Eastern States of India, *Records of the Zoological Survey of India*, Misc. ace. Paper no. 64, ZSI, Calcutta, pp. 216.
50. Sen, N. 2000. Occurrence, distribution and Status of diverified fish fauna of NorthEast India. : *Fish Germplasm Inventory and Conservation in NorthEast India* Pooniah, A.G. and Sarkar, U.K.(eds), NBFGR, Lucknow.
51. Sugunan, V. V. and Sinha, M., 2001: Sustainable capture and culture-based fisheries of India. *In: Pandian, T.J. (ed), Sustainable Indian Fisheries*. National Academy of Agricultural Sciences, New Delhi, pp 43-70.
52. Sarkar, U. K. and Ponniah, A. G. 2000. Evaluation of North East Indian fishes for their potential as cultivable, sport and ornamental fishes along with their conservation and endemic status, pp. 11-30. *In: Fish Biodiversity of North East India (eds. Ponniah, A.G. and Sarkar, U.K.). NATPpubl. 2. NBFGR*, Lucknow, India.

53. Talwar, P. K. and Jhingran, V. G. 1991; Inland Fishes of India and Adjacent Countries, vol. I & II, Oxford and IBH Publ. Co. Pvt. Ltd., Delhi, pp. ix+1097.
54. Tamang, P. 1992. Systematics, Distribution and Ecology of the ichthyospecies of Sikkim and their bearing on the fish and fisheries of the state, PhD Thesis, Gauhati University, pp. x+401.
55. Talwar, P. K. and Jhingran, V. G. 1991. Inland Fishes of India and Adjacent Countries, vol. I & II, Oxford and IBH Publ. Co. Pvt. Ltd., Delhi, pp. ix+1097.
56. Viswanath, W. and Tombi, H. 1987: On a collection of some endemic fishes of Manipur. Proc. Nat. Acad. Sci. 578, 11: 109.
57. Yadava, Y. S. 1987b. Studies on the ecology of an oxbow lake in context to development of beel in Assam. In: Workshop on Development of Beel Fishery in Assam. 21st and 22nd April, 1987, Assam Agricultural University, Jorhat, pp. 70-94.

